

P O L I S H J O U R N A L O F E N T O M O L O G Y
P O L S K I E P I S M O E N T O M O L O G I C Z N E

VOL. 78: 177-184

Bydgoszcz

30 June 2009

**Checklist of soil Mesostigmata (Acari) of Central Croatia (Dalmatia)
with some microenvironmental remarks**

SŁAWOMIR KACZMAREK *, TOMASZ MARQUARDT,
KATARZYNA FALEŃCZYK-KOZIRÓG

Kazimierz Wielki University, Institute of Environmental Biology
Department of Zoology, Ossolińskich Av. 12, PL-85-094 Bydgoszcz, Poland
*e-mail: slawkacz@ukw.edu.pl

ABSTRACT. Mesostigmatid mites (Acari) were studied in various microhabitats of Central Croatia (Dalmatia). Altogether 320 samples (taken during August 2002) and 2074 individuals were examined. Checklist of soil Mesostigmata, with microenvironmental notices was prepared.

KEY WORDS: Checklist, Acari, Mesostigmata, Croatia, Dalmatia, microenvironments.

INTRODUCTION

Croatia, because of its location within the Balkan Peninsula, is very interesting region of ecological investigations. Country has relatively small area with diversified climate and landscape which favours diversity of micro-environments and the formation of species richness. So far, the mite fauna of Croatia had been recognised only partially (mostly as Oribatida, Hydrachnidia and Uropodina) (HIRSCHMANN & WIŚNIEWSKI 1993, RUCNER & RUCNER 1995, PESIC 2002).

Study area

Soil samples were taken from various microhabitats within the area of Krka National Park, Brač Island and surroundings of a place called Makarska.

MATERIAL AND METHODS

Altogether 320 soil samples (50 cm^3 each) were taken during August 2002 from various micro-environments (moss covering soil, stones and tree trunks as well as decay and litter). Microhabitats of Krka NP were represented by 93 samples. From Brač Island and Makarska, respectively, 94 and 133 samples were taken. Totally 2074 specimens (1020 adult stages and 1054 juveniles) were obtained during 6-days extraction in Tullgren funnels. Mites were preserved in 70% ethyl alcohol and mounted in Hoyer's medium. Mites from order Mesostigmata (both mature and immature specimens), were identified to species level using universally applied keys (MICHERDZIŃSKI 1969, BŁASZAK 1974, BREGETOVA 1977, KARG 1993). Overall, 45 species of Mesostigmata (17 families) were recorded. Presented classification of Mesostigmata follows Joel Hallan's Biology Catalog (HALLAN 2005).

Checklist of Central Croatia (Dalmatia) soil Mesostigmata

Suborder: Epicriina

Superfamily: Epicriioidea

Family: Zerconidae BERLESE, 1892

Zercon athiasi VINCZE, 1965 – 4 ad. and 1 juv.

Brač – 1 ad. (moss covering stones); 1 ad. (litter)

Makarska – 1 ad. (moss covering soil); 1 ad. and 1 juv. (moss on tree trunks)

Zercon berlesei SELLNICK, 1958 – 2 ad. and 5 juv.

Brač – 2 ad. and 4 juv. (litter)

Krka NP – 1 juv. (moss covering stones)

Zercon fageticola HALAŠKOVÁ, 1969 – 32 ad. and 19 juv.

Makarska – 7 ad. and 5 juv. (moss covering soil); 18 ad. and 5 juv. (moss on stones);

5 ad. and 8 juv. (moss on tree trunks); 1 ad. (decay); 1 ad. and 1 juv. (litter)

Suborder: Uropodina

Superfamily: Uropodoidea

Family: Polyaspididae BERLESE, 1913

Polyaspis patavinus BERLESE, 1881 – 170 ad. and 429 juv.

Brač – 170 ad. and 429 juv. (decay)

Family: Trachytidae TRAGÄRDH, 1938

Trachytes lambda BERLESE, 1903 – 17 ad. and 16 juv.

Krka NP – 2 ad. (decay); 15 ad. and 16 juv. (litter)

Family: Trematuridae BERLESE, 1917

Trichouropoda ovalis (C.L. KOCH, 1839) – 13 ad. and 16 juv.

Krka NP – 1 ad. (moss covering soil); 11 ad. and 16 juv. (decay); 1 ad. (litter)

Family: Urodinychidae BERLESE, 1917

Dinychus inermis (C.L. KOCH, 1841) – 37 ad. and 14 juv.

Krka NP – 12 ad. and 3 juv. (moss covering soil); 25 ad. and 11 juv. (moss on tree trunks)

Urobovella vinicolora (VITZTHUM, 1926) – 34 ad. and 9 juv.

Krka NP – 34 ad. and 9 juv. (decay)

Family: Uropodidae KRAMER, 1881

Uropoda minima KRAMER, 1882 – 3 ad.

Krka NP – 1 ad. (decay); 2 ad. (litter)

Suborder: Parasitina

Superfamily: Parasitoidea

Family: Parasitidae OUDEMANS, 1901

Paragamasus runciger (BERLESE, 1903) – 22 ad. and 29 juv.

Krka NP – 3 ad. and 3 juv. (moss on soil); 10 ad. and 11 juv. (moss covering stones); 9 ad. and 15 juv. (litter)

Parasitus hyalinus (WILLMANN, 1949) – 5 ad. and 21 juv.

Brač - 2 juv. (moss on stones); 2 ad. and 2 juv. (decay); 3 ad. and 17 juv. (litter)

Pergamasus crassipes (LINNAE, 1758) – 8 ad. and 4 juv.

Krka NP – 3 ad. and 1 juv. (moss covering soil); 3 ad. and 3 juv. (moss on stones); 1 ad. (moss on tree trunks)

Makarska – 1 ad. (moss covering stones)

Pergamasus septentrionalis (OUDEMANS, 1902) – 1 ad.

Krka NP – 1 ad. (moss on tree trunks)

Suborder: Dermanyssina

Superfamily: Ascoidea

Family: Ascidae OUDEMANS, 1905

Asca aphidioides (LINNAE, 1758) – 68 ad. and 26 juv.

Krka NP – 27 ad. and 22 juv. (litter); 21 ad. and 2 juv. (moss on stone)

Makarska – 3 ad. and 2 juv. (decay); 9 ad. (moss covering soil); 2 ad. (moss on tree trunks); 6 ad. (litter).

Asca nova WILLMANN, 1939 – 64 ad. and 26 juv.

Brač – 13 ad. and 4 juv. (moss on stones); 1 ad. and 1 juv. (decay); 22 ad. and 18 juv. (litter)

Makarska – 18 ad. and 1 juv. (moss covering soil); 8 ad. (moss on stones); 2 ad. and 2 juv. (moss on tree trunk)

Cheiroleius serratus (HALBERT, 1915) – 154 ad. and 65 juv.

Krka NP – 23 ad. and 6 juv. (moss covering soil); 34 ad. and 15 juv. (moss on stones); 96 ad. and 40 juv. (moss on tree trunks)

Makarska – 1 juv. (moss covering soil)

Brač – 1 ad. and 3 juv. (litter)

Gamasellodes bicolor (BERLESE, 1918) – 45 ad. and 49 juv.

Brač – 1 ad. (moss covering stones); 9 ad. and 15 juv. (decay); 10 ad. and 2 juv. (litter)

Makarska – 4 juv. (moss on tree trunks); 7 ad. (decay);

Krka NP – 3 ad. and 11 juv. (decay); 15 ad. and 17 juv. (litter)

Lastoseius finetorum KARG, 1971 – 9 ad. and 28 juv.

Krka NP – 9 ad. nad 28 juv. (litter)

Platyseius italicus (BERLESE, 1905) – 1 ad.

Brač – 1 ad. (litter)

Proctolaelaps pygmaeus (MÜLLER, 1860) – 6 ad.

Makarska – 3 ad. (moss covering soil); 1 ad. (moss on stones); 1 ad. (moss on tree trunks); 1 ad. (litter)

Protogamasellus primitivus KARG, 1962 – 21 ad. and 5 juv.

Brač – 10 ad. and 1 juv. (moss on stones); 2 ad. (decay); 6 ad. and 4 juv. (litter)

Krka NP – 1 ad. (decay)

Makarska – 2 ad. (moss covering soil)

Zercoseius spathuliger (LEONARDI, 1899) – 12 ad.

Brač – 12 ad. (litter)

Family: Phytoseiidae BERLESE, 1916

Amblyseius obtusus (C.L. KOCH, 1839) – 2 ad. and 8 juv.

Krka NP – 1 ad. (moss on soil), 1 ad. and 8 juv. (litter)

Family: Podocinidae BERLESE, 1913

Podocinum pacificum BERLESE, 1896 – 1 juv.

Krka NP – 1 juv. (litter)

Superfamily: Dermanysssoidea

Family: Laelapidae BERLESE, 1892

Hypoaspis aculeifer (CANESTRINI, 1883) – 30 ad. and 59 juv.

Brač – 3 juv. (moss on stones); 21 ad. and 35 juv. (decay); 1 ad. and 1 juv. (litter);

Makarska – 1 ad. (moss on tree trunks); 1 juv. (decay)

Krka NP – 3 juv. (moss covering soil); 5 ad. and 13 juv. (moss on stones); 1 ad. (decay); 1 ad. and 3 juv. (litter)

Hypoaspis claviger (BERLESE, 1883) – 10 ad.

Brač – 1 ad. (moss on stones); 3 ad. (litter)

Makarska – 1 ad. (moss on stones); 1 ad. (litter)

Krka NP – 2 ad. (moss on stones); 1 ad. (decay); 1 ad. (litter)

Hypoaspis cuneifer (MICHAEL, 1891) – 1 ad.

Brač – 1 ad. (decay)

Hypoaspis kargi COSTA, 1968 – 34 ad. and 5 juv.

Krka NP – 5 ad. (moss covering stones); 5 ad. (litter)

Makarska – 5 ad. and 1 juv. (moss covering soil); 3 ad. (moss on stones); 3 ad. (moss on tree trunks); 1 ad. (decay); 3 ad. (litter)

Brač – 2 ad. and 1 juv. (moss on stones); 7 ad. and 3 juv. (litter);

Hypoaspis praesternalis WILLMANN, 1949 – 65 ad. and 75 juv.

Brač – 16 ad. and 25 juv. (moss on stones); 3 ad. and 3 juv. (decay); 22 ad. and 35 juv. (litter)

Makarska – 4 ad. (moss covering soil); 1 ad. (moss on tree trunks)

Krka NP – 1 ad. and 6 juv. (moss on stones); 9 ad. and 1 juv. (moss on tree trunks); 7 ad. and 4 juv. (decay); 2 ad. and 1 juv. (litter)

Hypoaspis vacua (MICHAEL, 1891) – 2 ad.

Makarska – 1 ad. (litter)

Brač – 1 ad. (litter)

Holostaspis montana (BERLESE, 1904) – 6 ad.

Makarska – 1 ad. (moss covering soil); 2 ad. (moss on stones)

Brač – 3 ad. (litter)

Pseudolaelaps paulseni (BERLESE, 1910) – 31 ad. and 5 juv.

Brač – 8 ad. and 5 juv. (moss covering stones); 6 ad. (decay); 3 ad. (litter)

Krka NP – 1 ad. (moss covering soil); 2 ad. (moss on stones); 4 ad. (decay)

Makarska – 5 ad. (moss covering soil); 1 ad. (moss on tree trunks); 1 ad. (litter)

Superfamily: Eviphiidoidea

Family: Eviphidiidae BERLESE, 1913

Evimirus uropodinus BERLESE, 1903 – 7 ad. and 4 juv.

Krka NP – 7 ad. and 3 juv. (moss covering stones); 1 juv. (decay)

Thinoseius kargi HIRSCHMANN, 1966 – 3 ad. and 1 juv.

Makarska – 1 juv. (moss covering soil); 1 ad. (moss on stones); 1 ad. (moss on tree trunks); 1 ad. (litter)

Family: Macrochelidae VITZTHUM, 1930

Macrocheles scutatiformis PETROVA, 1967 – 1 ad.

Krka NP – 1 ad. (moss on stones)

Macrocheles scutatus (BERLESE, 1904) – 3 ad. and 2 juv.

Krka NP – 2 ad. (moss covering soil); 1 ad. and 1 juv. (moss on stones); 1 juv. (decay)

Family: Pachylaelapidae BERLESE, 1913

Olopachys scutatus BERLESE, 1910 – 3 ad.

Makarska – 1 ad. (moss on soil); 2 ad. (litter)

Pachylaelaps pectinifer (G. et R. CANESTRINI, 1882) – 3 ad. and 7 juv.

Krka NP – 2 ad. and 3 juv. (moss on soil); 1 ad. and 2 juv. (moss covering stones); 2 juv. (decay)

Superfamily: Rhodacaroidea
Family: Digamasellidae EVANS, 1957

Dendrolaelaps rectus KARG, 1962 – 1 juv.
Krka NP – 1 juv. (moss covering stones)

Family: Rhodacaridae OUDEMANS, 1902

Rhodacarellus epigynalis SHEALS, 1956 – 27 ad. and 2 juv.
Brač – 13 ad. (moss covering stones); 14 ad. and 2 juv. (litter)
Rhodacarellus perspicuus HALAŠKOVÁ, 1958 – 25 ad. and 2 juv.
Brač – 20 ad. (moss covering stones); 5 ad. and 2 juv. (litter)
Rhodacarellus silesiacus WILLMANN, 1936 – 19 ad. and 112 juv.
Brač – 1 ad. (moss covering stones); 1 ad. (litter)
Krka NP – 1 ad. (moss covering stones); 2 ad. and 68 juv. (decay); 13 ad. and 44 juv.
(litter)
Makarska – 1 ad. (moss covering soil)

Superfamily: Veigaioidea
Family: Veigaiidae OUDEMANS, 1939

Veigaia cf. *decurtata* ATHIAS-HENRIOT, 1961 – 2 ad.
Makarska – 2 ad. (moss covering soil)
Veigaia exigua (BERLESE, 1916) – 10 ad. and 5 juv.
Brač – 4 ad. and 2 juv. (moss covering stones); 3 ad. and 3 juv. (litter)
Makarska – 3 ad. (litter)
Veigaia planicola (BERLESE, 1892) – 8 ad. and 3 juv.
Krka NP – 1 ad. and 1 juv. (moss covering stones); 3 ad. (decay); 4 ad. and 2 juv.
(litter)

REFERENCES

- BŁASZAK C. 1974. Zerconidae (Acari, Mesostigmata) Polski. The Zerconidae (Acari, Mesostigmata) of Poland. Monogr. Fauny Pol. 3: 315p.
- BREGETOVA I.G. 1977. Oprjedielitel obitajusczych w poczwje kleszczej. AN ZSSR: 5-718.
- HALLAN J. 2005. Synopsis of the described Mesostigmata of the world. Version 13 December 2008.
www site <http://insects.tamu.edu/research/collection/hallan/acari/> Mesostigmata1.htm (date of access 13 December 2008).

- HIRSCHMANN W., WIŚNIEWSKI J. 1993. Die Uropodiden der Erde. Acari Parasitiformes. Supercohors Atrichopygidiina, Hirschmann 1975. Acarologie Schriftenreihe für vergleichende Milbenkunde. Hirschman Verlag. Folge **40**. 466p.
- KARG W. 1993. Acari (Acarina), Milben Parasitiformes (Anactinochaeta) Cohors Gamasina Leach. Gustav Fischer Verlag. Jena. 523p.
- MICHERDZIŃSKI W. 1969. Die Familie Parasitidae Oudemans 1901 (Acarina, Mesostigmata). PWN, Warszawa: 666p.
- PESIC V.M. 2002. New records of water-mites (Acari, Hydrachnidia) based on the material collected by T. Petkovski in Croatia, including a check-list of species recorded in Croatia. Nat. Croat. Zagreb, vol. **11(4)**: 447-453.
- RUCNER D., RUCNER R. 1995. Contribution to the knowledge of some Arthropoda (Scorpiones, Pseudoscorpiones, Araneae, Acari, Diplopoda and Chilopoda) in forest communities of Croatia. Nat. Croat. Zagreb, vol. **4(4)**: 185-225.

Received: November 14, 2008

Accepted: April 17, 2009